



Development of a Unified SHOC-EDMF PBL Parameterization in the NASA GEOS Model

Nathan Arnold

Global Modeling and Assimilation Office,
NASA Goddard Space Flight Center

Kay Suselj and Joao Teixeira

NASA Jet Propulsion Laboratory,
California Institute of Technology


SHOC and EDMF background

Simplified HOC (Bogenschutz & Krueger 2013)

- Diagnoses higher order moments
- Analytic double gaussian for joint w , θ , q_t provides buoyancy flux to prognose TKE
- All fluxes downgradient

$$\overline{w'\theta'_l} = -K_H \frac{\partial \overline{\theta}_l}{\partial z}$$

$$\overline{\theta'^2_l} = \tau_v K_H \frac{\partial \overline{\theta}_l}{\partial z} \frac{\partial \overline{\theta}_l}{\partial z}$$




$$\frac{\partial e}{\partial t} = -\frac{\partial}{\partial z} \overline{w'e'} - w \frac{\partial e}{\partial z} + \frac{g}{\theta_v} \overline{w'\theta'_v} - \epsilon_e$$

Eddy Diffusivity - Mass Flux

(JPL version, Suselj et al 2021)


$$\overline{w'\phi'} \approx \underbrace{-K \frac{\partial \overline{\phi}}{\partial z}}_{\text{ED}} + \sum_i \underbrace{M_i (\phi_{ui} - \overline{\phi})}_{\text{MF}}$$

- Small eddy transport represented by diffusion;
- Nonlocal updraft transport by mass flux;
- JPL: multi-plume with stochastic entrainment




Recent work has combined EDMF and HOC approaches

Chinita et. al., 2022:
SHOC+MF in DoE SCREAM



Witte et. al., 2022:
CLUBB+MF in NCAR CESM



Improved profiles when MF tendencies included, vs. SHOC/CLUBB alone.

No direct contribution from MF to higher order moments.

SHOC-MF in the NASA GEOS model

Our implementation is unique in that we include Mass Flux contributions to all HOM calculations:

$$\overline{w'\theta'_l} = -K_H \frac{\partial \overline{\theta}_l}{\partial z} + \underbrace{M_u(\theta_{lu} - \overline{\theta}_l)}_{\text{MF}}$$

$$\overline{\theta_l'^2} = \tau_v \left[K_H \frac{\partial \overline{\theta}_l}{\partial z} - \underbrace{M_u(\theta_{lu} - \overline{\theta}_l)}_{\text{MF}} \right] \frac{\partial \overline{\theta}_l}{\partial z}$$

This allows ADG PDF to represent entire subgrid area **including updrafts**.

Unified cloud treatment.

$\overline{w'^3}$, $\overline{\theta_l'^3}$, $\overline{q_t'^3}$ are diagnosed directly from MF updrafts.

"Classic" SHOC estimates third moments following Canuto et al., 2001.



Our methodology


SHOC+MF implemented in NASA GEOS model

- GCM used for NWP (12km), S2S (50km), reanalysis (25km), GCRM (3km)
- Testing in SCM:
 - ARM97 continental diurnal CBL (Brown et al 2002)
 - BOMEX trade cumulus (Siebesma et al 2003)
 - DYCOMS II-RF01 Nocturnal non-precipitating stratocumulus (Zhu et al 2005)
 - Other cases (not shown)
 - $dz=25m$, $dt=225s$

Comparing with LES:

- System for Atmospheric Modeling (SAM v6.11.8)
- ARM/BOMEX: $dx/y=50m$, $dz=40m$, 12×12 km
- DYCOMS: $dx/y=25m$, $dz=5m$, 6.4×6.4 km


BOMEX mean profiles





Good agreement in θ_l , q_t . Cloud too few, too shallow.

Variances underestimated when MF contribution not included

BOMEX:






$W'hl'$ dominated by
MF in cloud layer



Without MF contribution, variances are underestimated

Third moments diagnosed from MF

BOMEX:






Mass flux qualitatively captures third moments, with some discrepancies




SHOC-MF ADG joint PDFs compared with MF, LES

BOMEX:

z=400m




z=700m




Reasonable agreement between ADG PDF plume 1 and MF properties, and with LES.

DYCOMS II-RF01 mean profiles




Good agreement in θ_l , q_t , mixed layer depth. Condensate somewhat underestimated.

ARM SGP mean profiles



SHOC-MF captures the diurnal evolution of θ_l and $w'\theta_l'$

ARM SGP cloud evolution vs. LES



Good agreement in timing, cloud base, fraction and condensate. Cloud top too shallow.

Conclusions

- We have implemented a SHOC+MF parameterization in the NASA GEOS model;
- The approach is unique in that MF contributions are included in higher order moments determining the double-gaussian PDF;
- Including MF contributions yields more realistic (vs LES) HOMs compared with the SHOC-only approach, and allows the PDF to represent the updraft domain fraction;
- The scheme simulates well the BOMEX, DYCOMS, ARM97 cases;
- **Future Work** will include global testing and tuning for NWP, and potentially rewriting PDF code to relax assumptions and allow closer correspondence between PDF plume and updrafts.

